



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 445 646 A3

EUROPEAN PATENT APPLICATION

(21) Application number: 91102937.9

(51) Int. Cl.⁵: **C12N 15/53**, C07H 21/04,
C12P 13/02

(22) Date of filing: 27.02.91

(30) Priority: 28.02.90 JP 48078/90

(43) Date of publication of application:
11.09.91 Bulletin 91/37

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB IT LI NL SE

(88) Date of deferred publication of the search report:
08.01.92 Bulletin 92/02

(71) Applicant: **NITTO CHEMICAL INDUSTRY CO., LTD.**
No. 5-1, Marunouchi 1-chome Chiyoda-ku Tokyo(JP)

Applicant: **Beppu, Teruhiko**
No. 5-21, 1-chome
Horinouchi Suginami-ku, Tokyo(JP)

Applicant: **YAMADA, Hideaki**
19-1 Matsugasaki-Kinomoto-cho Sakyo-ku
Kyoto-shi Kyoto-Fu(JP)

(72) Inventor: **Teruhiko, Beppu**
5-21, Horinouchi 1-chome, Suginami-ku, Tokyo(JP)
Inventor: **Hideaki, Yamada**
19-1, Matsugasaki Kinomotocho, Sakyo-ku, Kyoto-shi, Kyoto-fu(JP)
Inventor: **Toru, Nagasawa**
1-7, Takano Higashihirakicho, Sakyo-ku, Kyoto-shi, Kyoto-fu(JP)
Inventor: **Sueharu Horinouchi**
3-16-403 Etsuchujima 1-chome, Koutou-ku, Tokyo(JP)
Inventor: **Makoto, Nishiyama**
16-11, Nishiochiai 2-chome, Shinjuku-ku, Tokyo(JP)

(74) Representative: **Vossius & Partner**
Siebertstrasse 4 P.O. Box 86 07 67
W-8000 München 86(DE)

(54) **DNA fragment encoding a polypeptide having nitrile hydratase activity, a transformant containing the DNA fragment and a process for the production of amides using the transformant.**

(57) The present invention discloses the amino acid sequence and nucleotide sequence of the α - and β -subunits of two types of nitrile hydratase derived from *Rhodococcus rhodochrous* J-1. The DNA fragment encoding nitrile hydratase is inserted into an expression vector and the recombinant vector is used for transformation. The transformant contains

multiple copies of the gene and can produce much higher levels of nitrile hydratase than conventionally used microorganisms.

EP 0 445-646-A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 91102937.9

DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. CL.5)
D,A	FR - A1 - 2 633 938 (TERUHIKO BEPPU et al.) * Claims * & JP-A-2-119 778 --	1,6-9	C 12 N 15/53 C 07 H 21/04 C 12 P 13/02
P,A	CHEMICAL ABSTRACTS, vol. 112, no. 13, March 26, 1990, Columbus, Ohio, USA O. IKEHATA et al. "Primary structure of nitrile hydratase deduced from the nucleotide sequence of a Rhodococcus species and its expression in Escherichia coli" page 176, right column, abstract-no. 112 972f & Eur. J. Biochem. 1989, 181 (3), 563-70 --	1	
D,A	EP - A2 - 0 307 926 (YAMADA et al.) * Abstract * ----	1,8,9	TECHNICAL FIELDS SEARCHED (Int. CL.5) C 12 N C 07 H C 12 P
The present search report has been drawn up for all claims			
Place of search VIENNA	Date of completion of the search 28-10-1991	Examiner WOLF	
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			
T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document			

19



Europäisches Patentamt
European Patent Office
Office européen des brevets



11 Publication number:

0 445 646 A2

12

EUROPEAN PATENT APPLICATION

21 Application number: 91102937.9

51 Int. Cl.⁵: **C12N 15/53**, C07H 21/04,
C12P 13/02

22 Date of filing: 27.02.91

The microorganism(s) has (have) been deposited with Fermentation Research Institute under numbers FERM BP-1478, BP-2777, BP-2778, BP-1937.

30 Priority: 28.02.90 JP 48078/90

43 Date of publication of application:
11.09.91 Bulletin 91/37

84 Designated Contracting States:
AT BE CH DE DK ES FR GB IT LI NL SE

71 Applicant: **NITTO CHEMICAL INDUSTRY CO., LTD.**
No. 5-1, Marunouchi 1-chome Chiyoda-ku
Tokyo(JP)

Applicant: Beppu, Teruhiko
No. 5-21, 1-chome
Horinouchi Suginami-ku, Tokyo(JP)

Applicant: YAMADA, Hideaki
19-1 Matsugasaki-Kinomoto-cho Sakyo-ku
Kyoto-shi Kyoto-Fu(JP)

72 Inventor: Teruhiko, Beppu
5-21, Horinouchi 1-chome, Suginami-ku,
Tokyo(JP)
Inventor: Hideaki, Yamada
19-1, Matsugasaki Kinomotocho, Sakyo-ku,
Kyoto-shi, Kyoto-fu(JP)
Inventor: Toru, Nagasawa
1-7, Takano Higashihirakicho, Sakyo-ku,
Kyoto-shi, Kyoto-fu(JP)
Inventor: Sueharu Horinouchi
3-16-403 Etsuchujima 1-chome, Koutou-ku,
Tokyo(JP)
Inventor: Makoto, Nishiyama
16-11, Nishiochiai 2-chome, Shinjuku-ku,
Tokyo(JP)

74 Representative: **Vossius & Partner**
Siebertstrasse 4 P.O. Box 86 07 67
W-8000 München 86(DE)

54 DNA fragment encoding a polypeptide having nitrile hydratase activity, a transformant containing the DNA fragment and a process for the production of amides using the transformant.

57 The present invention discloses the amino acid sequence and nucleotide sequence of the α - and β -subunits of two types of nitrile hydratase derived from *Rhodococcus rhodochrous* J-1. The DNA fragment encoding nitrile hydratase is inserted into an expression vector and the recombinant vector is used for transformation. The transformant contains multiple copies of the gene and can produce much higher levels of nitrile hydratase than conventionally used microorganisms.

EP 0 445 646 A2

The present invention relates to a DNA fragment derived from *Rhodococcus rhodochrous* J-1 and encoding a polypeptide having nitrile hydratase activity which hydrates nitriles to amides. The invention also relates to a recombinant DNA containing the above DNA fragment, and a transformant transformed with the recombinant DNA. The present invention further relates to a method of producing nitrile hydratase using the transformant and of amides using nitrile hydratase.

Nitrile hydratase or nitrilase is known as an enzyme that hydrates nitriles to amides. Microorganisms that produce nitrile hydratase include those belonging to the genus *Bacillus*, the genus *Bacteridium*, the genus *Micrococcus* and the genus *Brevibacterium* (See, JP-B-62-21517/1989, USP No. 4,001,081), the genus *Corynebacterium* and the genus *Nocardia* (See, JP-B-56-17918/1981, USP No. 4,248,968), the genus *Pseudomonas* (See, JP-B-59-37951/1984, USP No. 4,637,982), the genus *Rhodococcus*, the genus *Arthrobacter* and the genus *Microbacterium* (See, JP-A-61-162193/1986, EP-A-0188316), and *Rhodococcus rhodochrous* (See, JP-A-2-470/1990, EP-A-0307926).

Nitrile hydratase has been used to hydrate nitriles to amides. In the invention, microorganisms are engineered to contain multiple copies of a recombinant DNA encoding nitrile hydratase according to a recombinant DNA technology. The recombinant produces a remarkably high level of nitrile hydratase compared with conventionally used microorganisms.

The present inventors previously disclosed a DNA fragment derived from *Rhodococcus* sp. N-774 (FERM BP-1936) which also encodes a polypeptide having nitrile hydratase activity (JP-A-2-119778/1988).

In contrast, the present inventors utilizes a DNA fragment derived from *Rhodococcus rhodochrous* J-1 for the production of nitrile hydratase. We isolated the gene encoding nitrile hydratase, inserted the gene into a suitable plasmid vector and transformed an appropriate host with the recombinant plasmid, thus successfully obtained the transformant producing nitrile hydratase which has high activity also on aromatic nitriles.

The present invention relates to

- (1) a DNA^(H) fragment encoding a polypeptide having nitrile hydratase activity, said polypeptide comprising the $\alpha^{(H)}$ -subunit as defined in the Sequence Listing by SEQ ID: No. 1 and the $\beta^{(H)}$ -subunit as defined in the Sequence Listing by SEQ ID: No. 2;
- (2) a DNA^(L) fragment encoding a polypeptide having nitrile hydratase activity, said polypeptide comprising the $\alpha^{(L)}$ -subunit as defined in the Sequence Listing by SEQ ID: No. 3 and the $\beta^{(L)}$ -subunit as defined in the Sequence Listing by SEQ ID: No. 4;
- (3) the DNA^(H) fragment of (1) which contains a nucleotide sequence encoding said $\alpha^{(H)}$ - and $\beta^{(H)}$ -subunits, comprising the DNA sequence of the $\alpha^{(H)}$ -subunit as defined in the Sequence Listing by SEQ ID: No. 5 and the DNA sequence of the $\beta^{(H)}$ -subunit as defined in the Sequence Listing by SEQ ID: No. 6;
- (4) the DNA^(L) fragment of (2) which contains a nucleotide sequence encoding said $\alpha^{(L)}$ - and $\beta^{(L)}$ -subunits, comprising the DNA sequence of the $\alpha^{(L)}$ -subunit as defined in the Sequence Listing by SEQ ID: No. 7 and the DNA sequence of the $\beta^{(L)}$ -subunit as defined in the Sequence Listing by SEQ ID: No. 8;
- (5) a recombinant DNA comprising the DNA^(H) or the DNA^(L) of (1)-(4) in a vector;
- (6) a transformant transformed with the recombinant DNA of (5);
- (7) a method for the production of nitrile hydratase which comprises culturing the transformant as described in (6) and recovering nitrile hydratase from the culture;
- (8) a method for the production of amides which comprises hydrating nitriles using nitrile hydratase as described in (7) to form amides; and
- (9) a method for the production of amides which comprises culturing the transformant as described in (6), and hydrating nitriles using the resultant culture, isolated bacterial cells, treated matter thereof, or a fixed material of them, to form amides.

The present invention is described in detail as follows.

The present invention is carried out by the steps (1)-(8):

(1) Isolation and Purification of Nitrile Hydratase and Partial Amino Acid Sequencing of Nitrile Hydratase

Two types of nitrile hydratase (designated as H type and L type, respectively) are isolated and purified from *Rhodococcus rhodochrous* J-1 (FERM BP-1478) and the both enzymes are separated into α and β subunits using HPLC. N-Terminal amino acid sequence each of the subunits is determined and shown in the Sequence Listing by SEQ ID: Nos. 9-12.

(2) Preparation of a DNA Probe for a Nitrile Hydratase Gene

A DNA probe is prepared from JM105/pYUK121 (FERM BP-1937) as described in JP-A-2-119778/1990 due to the high degree of homology in the amino acid sequence between the nitrile hydratase β subunit of *Rhodococcus* sp. N-774 described in said Japanese Patent Official Gazette and those of *Rhodococcus rhodochrous* J-1. Plasmid pYUK121 containing nitrile hydratase gene derived from *Rhodococcus* sp. N-774 is prepared from a JM105/pYUK121 culture. pYUK121 DNA is digested with SphI and SalI. The SphI-SalI fragment contains the nitrile hydratase gene (shown in the Sequence Listing by SEQ ID: No. 13) of *Rhodococcus* sp. N-774. The DNA fragment is radiolabeled.

(3) Detection of a DNA Segment Containing a Nitrile Hydratase Gene from the Chromosome of *Rhodococcus rhodochrous* J-1

Chromosomal DNA is prepared from a *Rhodococcus rhodochrous* J-1 culture. The chromosomal DNA is digested with restriction enzymes and hybridized to the probe described in (2) using the Southern hybridization method [Southern, E.M., J. Mol. Biol. 98, 503 (1975)].

Two DNA fragments of a different length are screened.

(4) Construction of a Recombinant Plasmid

A recombinant plasmid is constructed by inserting the chromosomal DNA fragment as prepared in (3) into a plasmid vector.

(5) Transformation and Screening for a Transformant Containing the Recombinant Plasmid

Transformants are prepared using the recombinant plasmid as described in (4). The transformant containing the recombinant plasmid is selected using the probe as described in (2) according to the colony hybridization method [R. Bruce Wallace et. al., Nuc. Ac. Res. 9, 879 (1981)]. Additionally, the presence of the nitrile hydratase gene in the recombinant plasmid is confirmed using the Southern hybridization method. The plasmids thus selected are designated as pNHJ10H and pNHJ20L.

(6) Isolation and Purification of Plasmid DNA and Construction of the Restriction Map

Plasmid DNAs of pNHJ10H and pNHJ20L as prepared in (5) are isolated and purified. The restriction map of the DNAs is constructed (Fig. 1) to determine the region containing nitrile hydratase gene.

(7) DNA Sequencing

The extra segment of the inserted DNA fragment in pNHJ10H and pNHJ20L is excised using an appropriate restriction enzyme. The inserted DNA fragment is then used for sequencing. The nucleotide sequence of the DNA fragment (SEQ: ID Nos. 14, 15) reveals that it contains the sequence deduced from the amino acid sequence as described in (1).

(8) Production of Nitrile Hydratase Using the Transformant and Conversion of Nitriles to Amides

The transformant as described in (8) is cultured. The bacterial cells are mixed with nitriles, a substrate of nitrile hydratase, and amides are produced.

Rhodococcus rhodochrous J-1 was deposited with Fermentation Research Institute, Agency of Industrial Science and Technology, and was assigned the accession number FERM BP-1478. A transformant TGI/pNHJ10H containing pNHJ10H as described in (5) and a transformant TGI/pNHJ20L containing pNHJ20L as described in (5) were deposited with the above and assigned the accession number FERM BP-2777 and FERM BP-2778, respectively.

Any vectors including a plasmid vector (e.g., pAT153, pMP9, pHC624, pKC7, etc.), a phage vector (e.g., λ gt11 (Toyobo), Charon 4A (Amersham), etc.) may be used. Enzymes which may be used include SphI, SalI, EcoRI, BamHI, SacI, and the like, which are commercially available (Takara Shuzo). Various hosts may be used for transformation including but not limited to *E. coli* JM105 and *E. coli* TGI.

Culture media for the transformant are those ordinarily used in the art.

Conversion of nitriles to amides is carried out using nitrile hydratase, crude nitrile hydratase, the culture of the transformant, the isolated bacterial cells or treated matter thereof, and the like, prepared from the culture of the transformant.

Suitable nitriles in the invention include aromatic nitriles having 4-10 carbon atoms in the aromatic moiety and aliphatic nitriles having 2-6 carbon atoms, which are described in the European Patent Publication No. 0,307,926. Typical examples of the nitriles are 4-, 3- and 2-cyanopyridines, benzonitrile, 2,6-difluorobenzonitrile, 2-thiophene carbonitrile, 2-furonitrile, cyanopyrazine, acrylonitrile, methacrylonitrile, crotonitrile, acetonitrile and 3-hydroxypropionitrile.

The present invention discloses the amino acid sequence and nucleotide sequence of the α - and β -subunits of two types of nitrile hydratase derived from *Rhodococcus rhodochrous* J-1. The DNA fragment encoding nitrile hydratase is inserted into an expression vector and the recombinant vector is used for transformation. The transformant contains multiple copies of the gene and can produce much higher levels of nitrile hydratase than conventionally used microorganisms.

Fig. 1 shows restriction maps of recombinant plasmids, pNHJ10H and pNHJ20L.

The present invention is illustrated by the following Example.

The following abbreviations are used in the Example.

TE:	Tris-HCl (10 mM; pH 7.8), EDTA (1 mM, pH 8.0)
TNE:	Tris-HCl (50 mM; pH 8.0), EDTA (1 mM, pH 8.0), NaCl (50 mM)
STE:	Tris-HCl (50 mM; pH 8.0), EDTA (5 mM, pH 8.0), Sucrose (35 mM)
2xYT medium:	1.6% Trypton; 1.0% Yeast extract, 0.5% NaCl

Example

(1) Isolation and Purification of Nitrile Hydratase and Partial Amino Acid Sequencing of Nitrile Hydratase

Rhodococcus rhodochrous J-1 was cultured in a medium (3 g/l of yeast extract, 0.5 g/l of KH_2PO_4 , 0.5 g/l of K_2HPO_4 , 0.5 g/l of $\text{MgSO}_4 \cdot 4\text{H}_2\text{O}$, 0.01 g/l of CoCl_2 , and 3 g/l of crotonamide, pH 7.2) at 28 °C for 80 hours. The bacterial cells were harvested. 50 g of the bacterial cells was disrupted and fractionated with ammonium sulfate. The sample was dialyzed and the dialysate was centrifuged. The supernatant was loaded on DEAE-Cellulofine chromatography, Phenyl-Sepharose chromatography, Sephadex G-150 chromatography and Octyl-Sepharose chromatography. Two fractions with enzyme activity were obtained and dialyzed. The dialysates were loaded on a high performance liquid chromatography using a reversed phase column (Senshu Pak VP-304-1251, Senshu Kagaku), and two respective subunits (α and β) were obtained. N-terminal amino acid sequence of $\alpha_1^{(H)}$, $\beta_1^{(H)}$, $\alpha_1^{(L)}$ and $\beta_1^{(L)}$ -subunits was determined using an Applied Biosystems model 470A protein sequencer. The amino acid sequences are shown in the Sequence Listing by SEQ ID: Nos. 9-12.

(2) Preparation of a DNA Probe for Nitrile Hydratase Gene

E. coli JM105 (FERM BP-1937) containing pYUK121 was cultured in 100 ml of 2xYT medium containing 50 $\mu\text{g}/\text{ml}$ of ampicillin at 30 °C overnight (12 hours). The bacterial cells were harvested and TNE was added to the cells. The cell suspension was then centrifuged. 8 ml of STE and 10 mg of lysozyme were added to the pellet. The mixture was incubated at 0 °C for five minutes followed by the addition of 4 ml of 0.25M EDTA. 2 ml of 10% SDS and 5 ml of 5M NaCl were then added to the mixture at room temperature. The resultant mixture was incubated at 0-4 °C for three hours and then ultracentrifuged. 1/2 volume of 30% PEG 6000 was added to the supernatant. The mixture was incubated at 0-4 °C overnight (12 hours) and centrifuged. TNE was added to the pellet to bring the volume to 7.5 ml and CsCl was then added to the suspension. The mixture was centrifuged to remove proteins. Then, 300-500 mg/ml of ethidium bromide was added to the supernatant. The mixture was transferred to a centrifuge tube. The tube was heat-sealed and then ultracentrifuged. cccDNA was extracted using a peristaltic pump. A bit more than equal amount of isopropyl alcohol saturated with water was added to the extract to rid of ethidium bromide. The sample was dialyzed against TE. About 3 ml of purified pYUK121 was obtained.

pYUK121 DNA was digested with SphI and SalI, resulting in a 2.07 kb DNA fragment containing a nitrile hydratase gene derived from *Rhodococcus* sp. N-774. The fragment was radiolabeled with ^{32}P to produce a probe. The nucleotide sequence of the probe is shown in the Sequence Listing by SEQ ID: No. 13.

(3) Preparation of a DNA Fragment Containing a Nitrile Hydratase Gene of Chromosome

Rhodococcus rhodochrous J-1 was cultured in 100 ml of a medium (10 g/l of glucose, 0.5 g/l of KH_2PO_4 , 0.5 g/l of K_2HPO_4 , 0.5 g/l of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 1 g/l of yeast extract, 7.5 g/l of peptone, 0.01 g/l of CoCl_2 , 7.5 g/l of urea, 1% glycine or 0.2 $\mu\text{g}/\text{ml}$ of ampicillin, 1 l of water, pH 7.2). The bacterial cells

were harvested and the pellet was washed with TNE. The pellet was then suspended in 10 ml of TE. 4 ml of 0.25M EDTA, 10-20 mg of lysozyme, 10-20 mg of achromoprotease and 10 ml of 10×SDS were added to the suspension. The suspension was incubated at 37°C for three hours. 15 ml of phenol was added to the suspension. The mixture was incubated at room temperature for 15 minutes and then centrifuged. The upper layer was removed, and 0.7 ml of 2.5M sodium acetate and diethyl ether were added to the supernatant. The mixture was centrifuged and the upper layer was discarded. Two volumes of ethanol were added to the bottom layer and DNA was removed with a glass rod. DNA was rinsed for five minutes each with TE:ethanol 2:8, 1:9, and 0:10 (v/v). DNA was then resuspended in 2-4 ml of TE (37°C). 10 µl of a mixture of RNase A and T₁ was added to the suspension and the mixture was incubated at 37°C. An equal amount of phenol was added to the mixture which was then centrifuged. More than equal amount of ether was added to the supernatant. The mixture was centrifuged again, and the upper layer was discarded and the bottom layer was saved. The bottom layer was dialyzed against 2 l of TE containing a small amount of chloroform overnight and further dialyzed against fresh TE for 3-4 hours. 4 ml of crude chromosomal DNA was obtained.

10 µl of TE, 3 µl of reaction buffer (10×) and 2 µl of SacI were added to 15 µl of crude chromosomal DNA. The mixture was incubated at 37°C for an hour and electrophoresed on an agarose gel at 60 V for three hours. The Southern hybridization of chromosomal DNA was carried out using the probe as described in (2). About 6.0 kb and 9.4 kb fragments were found to show a strong hybridization.

15 µl of chromosomal DNA was digested with SacI and electrophoresed on an agarose gel, as described above. 6.0 kb and 9.4 kb DNA fragments were cut out from the gel and taken in three volumes each of 8M NaClO₄. After solubilization, each solution was dotted on GF/C (Whatman) filter paper (6 mm in diameter). Ten drops (≈ 100 µl) of TE containing 6M NaClO₄ and then ten drops (≈ 100 µl) of 95% ethanol were added to the filter paper. The paper was air-dried for 3 minutes and placed in 0.5 ml Eppendorf tube. 40 µl of TE was added to the tube and the whole was incubated at 47°C for 30 minutes. The tube was then centrifuged. About 40 µl of the supernatant was obtained which contained 6.0 kb and 9.4 kb DNA fragments containing a nitrile hydratase gene of chromosomal DNA.

The method of inserting the 6.0 kb DNA fragment into a vector is described below. The same method is applied for the insertion of the 9.4 kb DNA fragment into a vector.

(4) Insertion of the Chromosomal DNA Fragment into a Vector

10 µl of TE, 3 µl of reaction buffer (10×) and 2 µl of SacI was added to 10 µl of pUC19. The mixture was incubated at 30°C for an hour. 2 µl of 0.25M EDTA was added to the mixture to stop the reaction. Then, 7 µl of 1M Tris-HCl (pH 9) and 3 µl of BAP (bacterial alkaline phosphatase) were added to the mixture. The mixture was incubated at 65°C for an hour. TE was then added to the mixture to make a total volume to 100 µl. The mixture was extracted 3× with an equal amount of phenol. An equal amount of ether was added to the extract. The bottom layer was removed and 10 µl of 3M sodium acetate and 250 µl of ethanol were added to the bottom layer. The mixture was incubated at -80°C for 30 minutes, centrifuged, dried, and resuspended in TE.

5 µl of pUC19 DNA thus obtained and 40 µl of the 6.0 kb DNA fragment as described in (3) were mixed. 6 µl of ligation buffer, 6 µl of ATP (6 mg/ml) and 3 µl of T4 DNA ligase were added to the mixture. The mixture was incubated at 4°C overnight (12 hours) to produce the recombinant plasmid containing the 6.0 kb DNA fragment encoding the desired enzyme in the SacI site of pUC19.

(5) Transformation and Screening of Transformants

E. coli TGI (Amersham) was inoculated into 10 ml of 2×YT medium and incubated at 37°C for 12 hours. After incubation, the resultant culture was added to fresh 2×YT medium to a concentration of 1%, and the mixture was incubated at 37°C for two hours. The culture was centrifuged and the pellet was suspended in 5 ml of cold 50 mM CaCl₂. The suspension was placed on ice for 40 minutes and then centrifuged. 0.25 ml of cold 50 mM CaCl₂ and 60 µl of the recombinant DNA as described in (4) were added to the pellet. The mixture was incubated at 0°C for 40 minutes, heat-shocked at 42°C for two minutes, placed on ice for five minutes, and added to 10 ml of 2×YT medium. The mixture was incubated at 37°C for 90 minutes with shaking, then centrifuged. The pellet was suspended in 1 ml of 2×YT medium, and two 10 µl aliquots of the suspension were plated on a 2×YT agar plate containing 50 µg/ml of ampicillin separately. The plate was incubated at 37°C. The colony grown on the plate was selected by the colony hybridization method: The colony was transferred to a nitrocellulose filter and digested. The DNA was fixed on the filter and hybridized to the probe as described in (2). The filter was autoradiographed and

a recombinant colony was selected. Additionally, the presence of a nitrile hydratase gene in the transformant was confirmed according to the Southern hybridization method.

(6) Isolation and Purification of Recombinant Plasmid and Construction of the Restriction Map of the Inserted DNA Fragments

The transformant selected as described in (5) was grown in 100 ml of 2×YT medium containing 50 µg/ml of ampicillin at 37 °C overnight (12 hours). The bacterial cells were harvested and TNE was added to the cells. The cells were collected again by centrifugation, and 8 ml of STE and 10 mg of lysozyme were added to the cells. The mixture was incubated at 0 °C for five minutes. 4 ml of 0.25M EDTA, 2 ml of 10% SDS (at room temperature) and 5 ml of 5M NaCl were added to the mixture. The mixture was incubated at 0-4 °C for three hours, and ultracentrifuged. 1/2 volume of 30% PEG 6000 was added to the supernatant. The mixture was incubated at 0-4 °C overnight (12 hours) and centrifuged again. TNE was added to the pellet to bring the volume up to 7.5 ml. CsCl was added to the suspension to rid of proteins. Then, 300-500 mg/ml of ethidium bromide was added to the supernatant and the mixture was transferred to a centrifuge tube. The tube was heat-sealed and ultracentrifuged. cccDNA was removed using a peristaltic pump. A bit more than equal amount of isopropyl alcohol saturated with water was added to cccDNA to remove ethidium bromide. The DNA sample was dialyzed against TE, resulting in about 3 ml of purified recombinant DNA. The recombinant plasmid thus obtained containing a 6.7 kb DNA fragment was designated as pNHJ10H. (The recombinant plasmid containing a 9.4 kb DNA fragment was designated as pNHJ20L).

These plasmid DNAs were digested with EcoRI, BamHI, PstI, SacI and Sall. The restriction maps were constructed and are shown in Fig. 1.

(7) DNA Sequencing

The location of a nitrile hydratase gene in the DNA fragment of pNHJ10H was determined according to the restriction map constructed and to the Southern hybridization method. An extra segment in pNHJ10H was cleaved off with PstI and Sall: The 6.0 kb DNA fragment resulted in 1.97 kb. Similarly, an extra segment in pNHJ20L was cleaved off with EcoRI and SacI: The 9.4 kb DNA fragment resulted in 1.73 kb. These DNA fragments were sequenced by the Sanger method [Sanger, F., Science 214: 1205-1210 (1981)] using M13 phage vector. The nucleotide sequence of the 1.97 kb DNA fragment (pNHJ10H) and the 1.73 kb DNA fragment (pNHJ20L) are shown in the Sequence Listing by SEQ ID: No. 14 and SEQ ID: No. 15, respectively.

The amino acid sequence deduced from the nucleotide sequence was found fully identical to the amino acid sequence as determined in (1). The sequence analysis also revealed that the DNA fragment contained the sequence coding for the α- and β-subunits.

(8) Production of Nitrile Hydratase Using the Transformant and Conversion of Nitriles to Amides Using Nitrile Hydratase

TG1/pNHJ10H and TG1/pNHJ20L were inoculated into 10 ml of 2×YT medium containing 50 µg/ml of ampicillin and incubated at 30 °C overnight (12 hours). 1 ml of the resultant culture was added to 100 ml of 2×YT medium (50 µg/ml of ampicillin, 0.1 g of CoCl₂·6H₂O/l). The mixture was incubated at 30 °C for 4 hours. IPTG was added to the mixture to a final concentration of 1 mM. The mixture was incubated at 30 °C for 10 hours. After harvesting the cells, the cells were suspended in 5 ml of 0.1 M phosphate buffer (pH 7.5). The suspensions were disrupted by sonification for 5 min and centrifuged at 12,000 ×g for 30 min. The resulting supernatants were used for the enzyme assay. The enzyme assay was carried out in a reaction mixture (12 ml) containing 50 mM potassium phosphate buffer (pH 7.5), 6 mM benzonitrile and an appropriate amount of the enzyme. The reaction was carried out at 20 °C for 30 min and stopped by the addition of 0.2 ml 1 M HCl. The amount of benzamide formed in the reaction mixture was determined by HPLC. As a control, the mixture obtained by the same procedure as described above but from E. coli TG1 was used. The levels of nitrile hydratase activity in cell-free extracts of E. coli containing pNHJ10H and pNHJ20L were 1.75×10^{-3} and 6.99×10^{-3} units/mg, respectively, when cultured in 2×YT medium in the presence of CoCl₂ and IPTG. Benzamide was found in the reaction mixture of TG1/pNHJ10H and pNHJ20L, whereas no benzamide was found in the reaction mixture of TG1.

(1) INFORMATION FOR SEQ ID NO: 1

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 203 amino acids

(B) TYPE: Amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(vi) ORIGINAL SOURCE

(A) ORGANISM: Rhodococcus rhodochrous

(B) STRAIN: J-1 (FERM BP-1478)

(ix) FEATURES

(A) OTHER INFORMATION

 $\alpha^{(H)}$ -subunit

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1

```

      5      10      15
MetSerGluHisValAsnLysTyrThrGluTyrGluAlaArgThr
      20      25      30
LysAlaIleGluThrLeuLeuTyrGluArgGlyLeuIleThrPro
      35      40      45
AlaAlaValAspArgValValSerTyrTyrGluAsnGluIleGly
      50      55      60
ProMetGlyGlyAlaLysValValAlaLysSerTrpValAspPro
      65      70      75
GluTyrArgLysTrpLeuGluGluAspAlaThrAlaAlaMetAla
      80      85      90
SerLeuGlyTyrAlaGlyGluGlnAlaHisGlnIleSerAlaVal
      95     100     105
PheAsnAspSerGlnThrHisHisValValValCysThrLeuCys
     110     115     120
SerCysTyrProTrpProValLeuGlyLeuProProAlaTrpTyr
     125     130     135
LysSerMetGluTyrArgSerArgValValAlaAspProArgGly

```

140 145 150
 ValLeuLysArgAspPheGlyPheAspIleProAspGluValGlu
 155 160 165
 ValArgValTrpAspSerSerSerGluIleArgTyrIleValIle
 170 175 180
 ProGluArgProAlaGlyThrAspGlyTrpSerGluGluGluLeu
 185 190 195
 ThrLysLeuValSerArgAspSerMetIleGlyValSerAsnAla
 200
 LeuThrProGlnGluValIleVal

(2) INFORMATION FOR SEQ ID NO: 2

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 229 amino acids

(B) TYPE: Amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(vi) ORIGINAL SOURCE

(A) ORGANISM: Rhodococcus rhodochrous

(B) STRAIN: J-1 (FERM BP-1478)

(ix) FEATURES

(A) OTHER INFORMATION

$\beta^{(H)}$ -subunit

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2

5 10 15
 MetAspGlyIleHisAspThrGlyGlyMetThrGlyTyrGlyPro
 20 25 30
 ValProTyrGlnLysAspGluProPhePheHisTyrGluTrpGlu
 35 40 45
 GlyArgThrLeuSerIleLeuThrTrpMetHisLeuLysGlyIle
 50 55 60
 SerTrpTrpAspLysSerArgPhePheArgGluSerMetGlyAsn
 65 70 75
 GluAsnTyrValAsnGluIleArgAsnSerTyrTyrThrHisTrp

```

5      80      85      90
LeuSerAlaAlaGluArgIleLeuValAlaAspLysIleIleThr
      95      100      105
GluGluGluArgLysHisArgValGlnGluIleLeuGluGlyArg
      110      115      120
10    TyrThrAspArgLysProSerArgLysPheAspProAlaGlnIle
      125      130      135
GluLysAlaIleGluArgLeuHisGluProHisSerLeuAlaLeu
      140      145      150
ProGlyAlaGluProSerPheSerLeuGlyAspLysIleLysVal
      155      160      165
15    LysSerMetAsnProLeuGlyHisThrArgCysProLysTyrVal
      170      175      180
ArgAsnLysIleGlyGluIleValAlaTyrHisGlyCysGlnIle
      185      190      195
20    TyrProGluSerSerSerAlaGlyLeuGlyAspAspProArgPro
      200      205      210
LeuTyrThrValAlaPheSerAlaGlnGluLeuTrpGlyAspAsp
      215      220      225
25    GlyAsnGlyLysAspValValCysValAspLeuTrpGluProTyr
LeuIleSerAla

```

(3) INFORMATION FOR SEQ ID NO: 3

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 207 amino acids

(B) TYPE: Amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(vi) ORIGINAL SOURCE

(A) ORGANISM: *Rhodococcus rhodochrous*

(B) STRAIN: J-1 (FERM BP-1478)

(ix) FEATURES

(A) OTHER INFORMATION

 $\alpha^{(L)}$ -subunit

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3

5 MetThrAlaHisAsnProValGlnGlyThrLeuProArgSerAsn
 10 GluGluIleAlaAlaArgValLysAlaMetGluAlaIleLeuVal
 AspLysGlyLeuIleSerThrAspAlaIleAspHisMetSerSer
 15 ValTyrGluAsnGluValGlyProGlnLeuGlyAlaLysIleVal
 AlaArgAlaTrpValAspProGluPheLysGlnArgLeuLeuThr
 AspAlaThrSerAlaCysArgGluMetGlyValGlyGlyMetGln
 20 GlyGluGluMetValValLeuGluAsnThrGlyThrValHisAsn
 MetValValCysThrLeuCysSerCysTyrProTrpProValLeu
 25 GlyLeuProProAsnTrpTyrLysTyrProAlaTyrArgAlaArg
 AlaValArgAspProArgGlyValLeuAlaGluPheGlyTyrThr
 ProAspProAspValGluIleArgIleTrpAspSerSerAlaGlu
 30 LeuArgTyrTrpValLeuProGlnArgProAlaGlyThrGluAsn
 PheThrGluGluGlnLeuAlaAspLeuValThrArgAspSerLeu
 35 IleGlyValSerValProThrThrProSerLysAla

(4) INFORMATION FOR SEQ ID NO: 4

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 226 amino acids

(B) TYPE: Amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(vi) ORIGINAL SOURCE

(A) ORGANISM: *Rhodococcus rhodochrous*

(B) STRAIN: J-1 (FERM BP-1478)

(ix) FEATURES

(A) OTHER INFORMATION

 β (L)-subunit

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4

```

      5      10      15
MetAspGlyIleHisAspLeuGlyGlyArgAlaGlyLeuGlyPro
      20      25      30
IleLysProGluSerAspGluProValPheHisSerAspTrpGlu
      35      40      45
ArgSerValLeuThrMetPheProAlaMetAlaLeuAlaGlyAla
      50      55      60
PheAsnLeuAspGlnPheArgGlyAlaMetGluGlnIleProPro
      65      70      75
HisAspTyrLeuThrSerGlnTyrTyrGluHisTrpMetHisAla
      80      85      90
MetIleHisHisGlyIleGluAlaGlyIlePheAspSerAspGlu
      95     100     105
LeuAspArgArgThrGlnTyrTyrMetAspHisProAspAspThr
     110     115     120
ThrProThrArgGlnAspProGlnLeuValGluThrIleSerGln
     125     130     135
LeuIleThrHisGlyAlaAspTyrArgArgProThrAspThrGlu
     140     145     150
AlaAlaPheAlaValGlyAspLysValIleValArgSerAspAla
     155     160     165
SerProAsnThrHisThrArgArgAlaGlyTyrValArgGlyArg
     170     175     180
ValGlyGluValValAlaThrHisGlyAlaTyrValPheProAsp
     185     190     195
ThrAsnAlaLeuGlyAlaGlyGluSerProGluHisLeuTyrThr
     200     205     210
ValArgPheSerAlaThrGluLeuTrpGlyGluProAlaAlaPro
     215     220     225
AsnValValAsnHisIleAspValPheGluProTyrLeuLeuPro
Ala

```

(5) INFORMATION FOR SEQ ID NO: 5

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 609 base pairs

(B) TYPE: Nucleic acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA

(vi) ORIGINAL SOURCE

(A) ORGANISM: Rhodococcus rhodochrous

(B) STRAIN: J-1 (FERM BP-1478)

(ix) FEATURES

(A) OTHER INFORMATION

 $\alpha^{(H)}$ -subunit

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5

```

      15      30      45
GTGAGCGAGCACGTCAATAAGTACACGGAGTACGAGGCACGTACC
      60      75      90
AAGGCGATCGAAACCTTGCTGTACGAGCGAGGGCTCATCAGCCC
     105     120     135
GCCGCGGTGCGACCGAGTCGTTTCGTACTACGAGAACGAGATCGGC
     150     165     180
CCGATGGGCGGTGCCAAGGTCGTGGCCAAGTCCTGGGTGGACCCT
     195     210     225
GAGTACCGCAAGTGGCTCGAAGAGGACGCGACGGCCGCGATGGCG
     240     255     270
TCATTGGGCTATGCCGGTGAGCAGGCACACCAAATTTGGCGGTC
     285     300     315
TTCAACGACTCCCAAACGCATCACGTGGTGGTGTGCACTCTGTGT
     330     345     360
TCGTGCTATCCGTGGCCGGTGCTTGGTCTCCCGCCCGCCTGGTAC
     375     390     405
AAGAGCATGGAGTACCGGTCCCGAGTGGTAGCGGACCCTCGTGGA
     420     435     450
GTGCTCAAGCGCGATTTCGGTTTCGACATCCCCGATGAGGTGGAG

```

5 GTCAGGGTTTGGGACAGCAGCTCCGAAATCCGCTACATCGTCATC
 CCGGAACGGCCGGCCGGCACCGACGGTTGGTCCGAGGAGGAGCTG
 10 ACGAAGCTGGTGAGCCGGGACTCGATGATCGGTGTCAGTAATGCG
 CTCACACCGCAGGAAGTGATCGTA

(6) INFORMATION FOR SEQ ID NO: 6

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 687 nucleic acids

(B) TYPE: Nucleic acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA

(vi) ORIGINAL SOURCE

(A) ORGANISM: Rhodococcus rhodochrous

(B) STRAIN: J-1 (FERM BP-1478)

(ix) FEATURES

(A) OTHER INFORMATION

$\beta^{(H)}$ -subunit

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6

45 ATGGATGGTATCCACGACACAGGCGGCATGACCGGATACGGACCG
 GTCCCTATCAGAAGGACGAGCCCTTCTTCCACTACGAGTGGGAG
 GGTCCGACCCGTGTCAATTCTGACTTGGATGCATCTCAAGGGCATA
 50 TCGTGGTGGGACAAGTCGCGGTTCTTCCGGGAGTCGATGGGGAAC
 GAAAACTACGTCAACGAGATTCCGAACTCGTACTACACCCACTGG
 55 CTGAGTGCGGCAGAACGTATCCTCGTCGCGGACAAGATCATCACC

```

5      285      300      315
GAAGAAGAGCGAAAGCACCGTGTGCAAGAGATCCTTGAGGGTCCG
      330      345      360
TACACGGACAGGAAGCCGTCGCGGAAGTTCGATCCGGCCCAGATC
      375      390      405
10     GAGAAGGCGATCGAACGGCTTCACGAGCCCCACTCCCTAGCGCTT
      420      435      450
CCAGGAGCGGAGCCGAGTTTCTCTCTCGGTGACAAGATCAAAGTG
      465      480      495
15     AAGAGTATGAACCCGCTGGGACACACACGGTGCCCGAAATATGTG
      510      525      540
CGGAACAAGATCGGGGAAATCGTCGCCTACCACGGCTGCCAGATC
      555      570      585
20     TATCCCGAGAGCAGCTCCGCCGGCCTCGGCGACGATCCTCGCCCG
      600      615      630
CTCTACACGGTTCGCGTTTTTCGCCCCAGGAACTGTGGGGCGACGAC
      645      660      675
GGAAACGGGAAAGACGTAGTGTGCGTCGATCTCTGGGAACCGTAC
25     CTGATCTCTGCG

```

(7) INFORMATION FOR SEQ ID NO: 7

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 621 base pairs

(B) TYPE: Nucleic acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA

(vi) ORIGINAL SOURCE

(A) ORGANISM: *Rhodococcus rhodochrous* J-1

(FERM BP-1478)

(ix) FEATURES

(A) OTHER INFORMATION

$\alpha^{(L)}$ -subunit

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7

5 ATGACCGCCCAACAATCCCGTCCAGGGCACGTTGCCACGATCGAAC
GAGGAGATCGCCGCACGCGTGAAGGCCATGGAGGCCATCCTCGTC
10 GACAAGGGCCTGATCTCCACCGACGCCATCGACCACATGTCTCTCG
GTCTACGAGAACGAGGTCTGCTCAACTCGGCGCCAAGATCGTC
GCCCGCGCCTGGGTCTGATCCCGAGTTCAAGCAGCGCCTGCTCACC
15 GACGCCACCAGCGCCTGCCGTGAAATGGGCGTGGGCGGCATGCAG
GGCGAAGAAATGGTCTGCTGGAAAACACCGGCACGGTCCACAAC
20 ATGGTCTGTATGTACCTTGTGCTCGTCTATCCGTGGCCGGTTCTC
GGCCTGCCACCCAACTGGTACAAGTACCCCGCCTACCGCGCCCGC
25 GCTGTCCGCGACCCCCGAGGTGTGCTGGCCGAATTCCGATATACC
CCCGACCCTGACGTCTGAGATCCGGATATGGGACTCGAGTGCCGAA
CTTCGCTACTGGGTCTCTGCCGCAACGCCAGCCGGCACCGAGAAC
30 TTCACCGAAGAACAACCTCGCCGACCTCGTCAACCGCGACTCGCTC
ATCGGCGTATCCGTCCCCACCACACCCAGCAAGGCC

(8) INFORMATION FOR SEQ ID NO: 8

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 678 base pairs

(B) TYPE: Nucleic acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA

(vi) ORIGINAL SOURCE

(A) ORGANISM: Rhodococcus rhodochrous

(B) STRAIN: J-1 (FERM BP-1478)

(ix) FEATURES

(A) OTHER INFORMATION

$\beta^{(L)}$ -subunit

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8

```

15      ATGGATGGAATCCACGACCTCGGTGGCCGCGCCGGCCTGGGTCCG
           15      30      45
20      ATCAAGCCCGAATCCGATGAACCTGTTTTCCATTCCGATTGGGAG
           60      75      90
25      CCGTCGGTTTTTGACGATGTTCCCGGCGATGGCGCTGGCCGGCGCG
           105     120     135
30      TTCAATCTCGACCAGTTCCGGGGCGCGATGGAGCAGATCCCCCG
           150     165     180
35      CACGACTACCTGACCTCGCAATACTACGAGCACTGGATGCACGCG
           195     210     225
40      ATGATCCACCACGGCATCGAGGCGGGCATCTTCGATTCCGACGAA
           240     255     270
45      CTCGACCGCCGCACCCAGTACTACATGGACCATCCGGACGACACG
           285     300     315
50      ACCCCCACGCGGCAGGATCCGCAACTGGTGGAGACGATCTCGCAA
           330     345     360
55      CTGATCACCACGGAGCCGATTACCGACGCCCGACCGACACCGAG
           375     390     405
60      GCCGCATTCCCGGTAGGCGACAAAGTCATCGTTCGGTCCGGACGCC
           420     435     450
65      TCACCGAACACCCACACCCGCCGCGCCGGATACGTCCGCGGTCTGT
           465     480     495
70      GTCGGCGAAGTCGTGGCGACCCACGGCGCGTATGTCTTTCCGGAC
           510     525     540
75      ACCAACGCACTCGGCGCCGGCGAAAGCCCCGAACACCTGTACACC
           555     570     585
80      GTGCGGTTCTCGGCGACCGAGTTGTGGGGTGAACCTGCCGCCCG
           600     615     630
85      AACGTCGTCAATCACATCGACGTGTTTGAACCGTATCTGCTACCG
           645     660     675
90      GCC

```

(9) INFORMATION FOR SEQ ID NO: 9

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 amino acids

(B) TYPE: Amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(vi) ORIGINAL SOURCE

(A) ORGANISM: Rhodococcus rhodochrous

(B) STRAIN: J-1 (FERM BP-1478)

(ix) FEATURES

(A) OTHER INFORMATION

$\alpha^{(H)}$ -subunit: $\alpha_1^{(H)}$

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9

Ser-Glu-His-Val-Asn-Lys-Tyr-Thr-Glu-Tyr-Glu-Ala-Arg-Thr-Lys
Ala-Ile-Glu-Thr-Leu-Leu-Tyr-Glu-Arg-Gly-Leu

(10) INFORMATION FOR SEQ ID NO: 10

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 amino acids

(B) TYPE: Amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(vi) ORIGINAL SOURCE

(A) ORGANISM: Rhodococcus rhodochrous

(B) STRAIN: J-1 (FERM BP-1478)

(ix) FEATURES

(A) OTHER INFORMATION

$\beta^{(H)}$ -subunit: $\beta_1^{(H)}$

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10

Met-Asp-Gly-Ile-His-Asp-Thr-Gly-Gly-Met-Thr-Gly-Tyr-Gly-Pro
Val-Pro-Tyr-Gln-Lys-Asp-Glu-Pro-Phe-Phe-His-Tyr-Glu

(11) INFORMATION FOR SEQ ID NO: 11

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 amino acids

(B) TYPE: Amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(vi) ORIGINAL SOURCE

(A) ORGANISM: Rhodococcus rhodochrous

(B) STRAIN: J-1 (FERM BP-1478)

(ix) FEATURES

(A) OTHER INFORMATION

$\alpha^{(L)}$ -subunit: $\alpha_1^{(L)}$

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11

Thr-Ala-His-Asn-Pro-Val-Gln-Gly-Thr-Leu-Pro-Arg-?-Asn-Glu

(12) INFORMATION FOR SEQ ID NO: 12

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 19 amino acids

(B) TYPE: Amino acid

(C) STRANDEDNESS:

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Peptide

(vi) ORIGINAL SOURCE

(A) ORGANISM: Rhodococcus rhodochrous

(B) STRAIN: J-1 (FERM BP-1478)

(ix) FEATURES

(A) OTHER INFORMATION

$\beta^{(L)}$ -subunit: $\beta_1^{(L)}$

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12

Met-Asp-Gly-Ile-His-Asp-Leu-Gly-Gly-Arg-Ala-?-Leu-?-Pro
Ile-Lys-Pro-Glu

(13) INFORMATION FOR SEQ ID NO: 13

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2070 base pairs
(B) TYPE: Nucleic acid
(C) STRANDEDNESS: Single
(D) TOPOLOGY: Linear
(ii) MOLECULE TYPE: Genomic DNA
(vi) ORIGINAL SOURCE
(A) ORGANISM: Rhodococcus sp.
(B) STRAIN: N-774 (FERM BP-1936)
(ix) FEATURES
from nucleotide No. 675 to 1295: subunit α
from nucleotide No. 1225 to 1960: subunit β
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13

SphI
GCATGCTTTCCACATCTGGAACGTGATCGCCACGGACGGTGGTG
CCTACCAGATGTTGGACGGCAACGGATACGGCATGAACGCCGAAG
GTTTGTACGATCCGGAAGTGTGGCACACTTTGCTTCTCGACGCA
TTCAGCACGCCGACGCTCTGTCCGAAACCGTCAAACCTGGTGGCCC
TGACCGGCCACACGGCATCACCCCTCGGCGGCGGAGCTACG
GCAAAGCCCGGAACCTCGTACCGCTTGCCCGCGCCGCTACGACA
CTGCCTTGAGACAATTCGACGCTCCTGGTGATGCCAACGCTGCCCT
ACGTCGCATCCGAATTGCCGGCGAAGGACGTAGATCGTGCAACCT

TCATCACCAAGGCTCTCGGGATGATCGCCAACACGGCACCATTTCG⁴⁰⁰
 5 ACGTGACCGGACATCCGTCCCTGTCCGTTCCGGCCGGCCTGGTGA
 .⁴⁵⁰
 ACGGGGTTCGGTTCGGAATGATGATCACCGGCAGACACTTCGACG
 10 ATGCGACAGTCCTTCGTGTCCGACGCGCATTTCGAAAAGCTTCGCG
 .⁵⁰⁰
 GCCGGTTTCCGACGCCGGCCGAACGCGCCTCCAACCTCTGCCACCAC
 .⁵⁵⁰
 AACTCAGCCCCCGCCTAGTCTCTGACGCACTGTCAGACAACAAATTC
 15 CACCGATTACACATGATCAGCCCACATAAGAAAAGGTGAACCAG
 .⁶⁰⁰
 ATGTCAGTAACGATCGACCACACAACGGAGAACGCCCGCACCGGCC
 20 MetSerValThrIleAspHisThrThrGluAsnAlaAlaProAla
 Subunit α
 CAGGCGGCGGTCTCCGACCGGGCGTGGGCACTGTTCCGCGCACTC
 .⁶⁵⁰
 GlnAlaAlaValSerAspArgAlaTrpAlaLeuPheArgAlaLeu
 Kpn I
 25 GACGGTAAGGGATTGGTACCCGACGGTTACGTCGAGGGATGGAAG
 AspGlyLysGlyLeuValProAspGlyTyrValGluGlyTrpLys
 .⁷⁰⁰
 AAGACCTCCGAGGAGGACTTCAGTCCAAGGCGCGGAGCGGAATTG
 30 LysThrSerGluGluAspPheSerProArgArgGlyAlaGluLeu
 Pvu II
 GTAGCGCGCGCATGGACCGACCCCGAGTTCCGGCAGCTGCTTCTC
 ValAlaArgAlaTrpThrAspProGluPheArgGlnLeuLeuLeu
 .⁷⁵⁰
 ACCGACGGTACCGCCGCAGTTGCCAGTACGGATACCTGGGCCCC
 35 ThrAspGlyThrAlaAlaValAlaGlnTyrGlyTyrLeuGlyPro
 .⁸⁰⁰
 CAGGCGGCCTACATCGTGGCAGTCGAAGACACCCCGACACTCAAG
 GlnAlaAlaTyrIleValAlaValGluAspThrProThrLeuLys
 40 AACGTGATCGTGTGCTCGCTGTGTTTCATGCACCGCGTGGCCCATC
 AsnValIleValCysSerLeuCysSerCysThrAlaTrpProIle
 .⁸⁵⁰
 CTCGGTCTGCCACCCACCTGGTACAAGAGCTTCGAATACCGTGCG
 45 LeuGlyLeuProProThrTrpTyrLysSerPheGluTyrArgAla

CGCGTGGTCCGCGAACCACGGAAGGTTCTCTCCGAGATGGGAACC
 ArgValValArgGluProArgLysValLeuSerGluMetGlyThr
 5 GAGATCGCGTCGGACATCGAGATTTCGCGTCTACGACACCACCGCC
 GluIleAlaSerAspIleGluIleArgValTyrAspThrThrAla
 GAAACTCGCTACATGGTCCTCCCGCAGCGTCCCGCCGGCACCAGAA
 10 GluThrArgTyrMetValLeuProGlnArgProAlaGlyThrGlu
 Pst I
 GGCTGGAGCCAGGAACAACCTGCAGGAAATCGTCACCAAGGACTGC
 GlyTrpSerGlnGluGlnLeuGlnGluIleValThrLysAspCys
 15 CTGATCGGGGTTGCAATCCCGCAGGTTCCCAACCGTCTGATCACCC
 LeuIleGlyValAlaIleProGlnValProThrValTRM
 CGACAAGAAGGAAGCACACC-ATGGATGGAGTACACGATCTTGCC
 20 MetAspGlyValHisAspLeuAla
 Subunit β
 GGAGTACAAGGCTTCGGCAAAGTCCCGCATACCGTCAACGCCGAC
 GlyValGlnGlyPheGlyLysValProHisThrValAsnAlaAsp
 25 ATCGGCCCCACCTTTTCACGCCGAATGGGAACACCTGCCCTACAGC
 IleGlyProThrPheHisAlaGluTrpGluHisLeuProTyrSer
 CTGATGTTCCCGGTGTCCCGAACTCGGGGCCTTCAGCGTCGAC
 30 LeuMetPheAlaGlyValAlaGluLeuGlyAlaPheSerValAsp
 Sal I
 GAAGTGCATACGTCTCGAGCGGATGGAGCCGGGCCACTACATG
 GluValArgTyrValValGluArgMetGluProGlyHisTyrMet
 35 ATGACCCCGTACTACGAGAGGTACGTCATCGGTGTGCGGACATTG
 MetThrProTyrTyrGluArgTyrValIleGlyValAlaThrLeu
 ATGGTCAAAAAGGGAATCCTGACGCAGGACGAACTCGAAAGCCTT
 40 MetValGluLysGlyIleLeuThrGlnAspGluLeuGluSerLeu
 GCGGGGGGACCGTTCCCACTGTACGGCCAGCGAATCCGAAGGG
 AlaGlyGlyProPheProLeuSerArgProSerGluSerGluGly
 45
 50
 55

CGGCCGGCACCCGTCGAGACGACCACCTTCGAAGTCGGGCAGCGA¹⁷⁰⁰
 ArgProAlaProValGluThrThrThrPheGluValGlyGlnArg
 5 GTACGCGTACGCGACGAGTACGTTCCGGGGCATATTCGAATGCCT¹⁷⁵⁰
 ValArgValArgAspGluTyrValProGlyHisIleArgMetPro
 GCATACTGCCGTGGACGAGTGGGAACCATCTCTCATCGAACTACC
 10 AlaTyrCysArgGlyArgValGlyThrIleSerHisArgThrThr
 GAGAAGTGGCCGTTTCCCGACGCAATCGGCCACGGGCGCAACGAC¹⁸⁰⁰
 GluLysTrpProPheProAspAlaIleGlyHisGlyArgAsnAsp
 GCCGGCGAAGAACCGACGTACCACGTGAAGTTCGCCGCCGAGGAA¹⁸⁵⁰
 AlaGlyGluGluProThrTyrHisValLysPheAlaAlaGluGlu
 TTGTTCCGGTAGCGACACCGACGGTGGAAGCGTCGTTGTGCGACCTC¹⁹⁰⁰ Sal I
 20 LeuPheGlySerAspThrAspGlyGlySerValValValAspLeu
 TTCGAGGGTTACCTCGAGCCTGCGGCCTGATCTTCCAGCATTCCA¹⁹⁵⁰
 PheGluGlyTyrLeuGluProAlaAlaTRM
 GGCGGCGGTCACGCGATCACAGCGGTTTCGTGCGACCGCCGCCTGA²⁰⁰⁰
 25 TCACCACGATTCACTCATTGGAAGGACACTGGAAATCATGGTCG²⁰⁵⁰
 Sal I
 AC

(14) INFORMATION FOR SEQ ID NO: 14

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1970 base pairs

(B) TYPE: Nucleic acid

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA

(vi) ORIGINAL SOURCE

(A) ORGANISM: *Rhodococcus rhodochrous* J-1

(FERM BP-1478)

(ix) FEATURES

from nucleotide No. 408 to 1094: subunit $\beta^{(H)}$ from nucleotide No. 1111 to 1719: subunit $\alpha^{(H)}$

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14

```

15      10      20      30      40      50      60
CTGCAGCTCGAACATCGAAGGGTGCGAGCCGAGAGATCGGAGACGCAGACACCCGGAGGG

20      70      80      90     100     110     120
AACTTAGCCTCCCGGACCGATGCGTGCTCTGGCAACGCCTCAAAATTCAGTGCAAGCGAT

25      130     140     150     160     170     180
TCAATCTTGTTACTTCCAGAACCGAATCACGTCCCCGTAGTGTGCGGGGAGAGCGCCCGA

30      190     200     210     220     230     240
ACGCAGGGGATGGTATCCATGCGCCCCCTTCTCTTTTCGAACGAGAACCGGCCGGTACAGCC

35      250     260     270     280     290     300
GACCCGGAGACACTGTGACGCCGTTCAACGATTGTTGTGCTGTGAAGGATTCACCCAAGC

30      310     320     330     340     350     360
CAACTGATATCGCCATTCCGTTGCCGGAACATTTGACACCTTCTCCCTACGAGTAGAAGC

35      370     380     390     400     410     420
CAGCTGGACCCCTCTTTGAGCCCAGCTCCGATGAAAGGAATGAGGAAATGGATGGTATCC
MetAspGlylleH
Subunit  $\beta^{(H)}$ 

40      430     440     450     460     470     480
ACGACACAGGCGGCATGACCGGATACGGACCGGTCCCCTATCAGAAGGACGAGCCCTTCT
isAspThrGlyGlyMetThrGlyTyrGlyProValProTyrGlnLysAspGluProPheP

45      490     500     510     520     530     540
TCCACTACGAGTGGGAGGGTCCGACCCTGTCAATTCTGACTTGGATGCATCTCAAGGGCA
heHisTyrGluTrpGluGlyArgThrLeuSerlleLeuThrTrpMetHisLeuLysGlyI

50      550     560     570     580     590     600
TATCGTGGTGGGACAAGTCGCGGTTCTTCCGGGAGTCGATGGGGAACGAAAACCTACGTCA
leSerTrpTrpAspLysSerArgPhePheArgGluSerMetGlyAsnGluAsnTyrValA

55      610     620     630     640     650     660
ACGAGATTTCGCAACTCGTACTACCCCACTGGCTGAGTGCGGCAGAACGTATCCTCGTCG
snGluIleArgAsnSerTyrTyrThrHisTrpLeuSerAlaAlaGluArglleLeuValA

60      670     680     690     700     710     720
CCGACAAGATCATCACCAGAAGAGCGAAAGCACCGTGTGCAAGAGATCCTTGAGGGTC
laAspLysllelleThrGluGluGluArgLysHisArgValGlnGluIleLeuGluGlyA

```

1450 1460 1470 1480 1490 1500
CCGGTGGCTTGGTCTCCCGCCCGCCTGGTACAAGAGCATGGAGTACCGGTCCCGAGTGGTA
ProValLeuGlyLeuProProAlaTrpTyrLysSerMetGluTyrArgSerArgValVal
5 1510 1520 1530 1540 1550 1560
GCGGACCCTCGTGGAGTGGTCAAGCGCGATTTTCGGTTTCGACATCCCGATGAGGTGGAG
AlaAspProArgGlyValLeuLysArgAspPheGlyPheAspIleProAspGluValGlu
10 1570 1580 1590 1600 1610 1620
GTCAGGGTTTGGGACAGCAGCTCCGAAATCCGCTACATCGTCATCCCGGAACGGCCGGCC
ValArgValTrpAspSerSerSerGluIleArgTyrIleValIleProGluArgProAla
1630 1640 1650 1660 1670 1680
GGCACCAGCGGTTGGTCCGAGGAGGAGCTGACGAAGCTGGTGAGCCGGGACTCGATGATC
GlyThrAspGlyTrpSerGluGluGluLeuThrLysLeuValSerArgAspSerMetIle
15 1690 1700 1710 1720 1730 1740
GGTGTGAGTAATGCGCTCACACCGCAGGAAGTGATCGTATGAGTGAAGACACACTCACTG
GlyValSerAsnAlaLeuThrProGlnGluValIleVal
1750 1760 1770 1780 1790 1800
ATCGGCTCCCGGCGACTGGGACCGCCGCACCGCCCCGCGACAATGGCGAGCTTGTATTCA
20 1810 1820 1830 1840 1850 1860
CCGAGCCTTGGGAAGCAACGGCATTCGGGGTCCGCCATCGCGCTTTCGGATCAGAAGTCGT
1870 1880 1890 1900 1910 1920
ACGAATGGGAGTTCTTCCGACAGCGTCTCATTCCTCCATCGCTGAGGCCAACGGTTGCG
25 1930 1940 1950 1960 1970
AGGCATACTACGAGAGCTGGACAAAGGCGCTCGAGGCCAGCGTGGTCGAC
30
35
40

(15) INFORMATION FOR SEQ ID NO: 15

- 45 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1731 base pairs
- 50 (B) TYPE: Nucleic acid
- 55

(C) STRANDEDNESS: Single

(D) TOPOLOGY: Linear

(ii) MOLECULE TYPE: Genomic DNA

(vi) ORIGINAL SOURCE

(A) ORGANISM: Rhodococcus rhodochrous

(B) STRAIN: J-1 (FERM BP-1478)

(ix) FEATURES

from nucleotide No. 171 to 848: subunit $\beta^{(L)}$

from nucleotide No. 915 to 1535: subunit $\alpha^{(L)}$

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15

```

      10      20      30      40      50      60
GAGCTCCCTGGAGCCACTCGCGCCGACGCATCCACGCTCGGACAGCCCACGGTGCGGATC

      70      80      90     100     110     120
ACCCCTGTTTCGTCGGTAACAGAACAGTAACATGTCATCAGGTCATGACGTGTTGACGCAT

      130     140     150     160     170     180
TAGACGAGGGCACATAGGGTTGGTGACTCACGGCACAAGGAGAGCATTTCATGGATGGAA
                               MetAspGlyI
                               Subunit  $\beta^{(L)}$ 

      190     200     210     220     230     240
TCCACGACCTCGGTGGCCGCGCCGGCCTGGGTCCGATCAAGCCCGAATCCGATGAACCTG
leHisAspLeuGlyGlyArgAlaGlyLeuGlyProIleLysProGluSerAspGluProV

      250     260     270     280     290     300
TTTTCCATTCCGATTGGGAGCGGTCGGTTTTTGACGATGTTCCCGGCGATGGCGCTGGCCG
alPheHisSerAspTrpGluArgSerValLeuThrMetPheProAlaMetAlaLeuAlaG

      310     320     330     340     350     360
GCGCGTTCAATCTCGACCAGTTCCGGGGCGCGATGGAGCAGATCCCCCGCAGCACTACC
lyAlaPheAsnLeuAspGlnPheArgGlyAlaMetGluGlnIleProProHisAspTyrL

      370     380     390     400     410     420
TGACCTCGCAATACTACGAGCACTGGATGCACGCGATGATCCACCACGGCATCGAGGCGG
euThrSerGlnTyrTyrGluHisTrpMetHisAlaMetIleHisHisGlyIleGluAlaG

      430     440     450     460     470     480
GCATCTTCGATTCCGACGAACCTCGACCGCGCACCCAGTACTACATGGACCATCCGGACG
lyIlePheAspSerAspGluLeuAspArgArgThrGlnTyrTyrMetAspHisProAspA

```

490 500 510 520 530 540
 ACACGACCCACGCGGCAGGATCCGCAACTGGTGGAGACGATCTCGCAACTGATCACCC
 spThrThrProThrArgGlnAspProGlnLeuValGluThrIleSerGlnLeuIleThrH
 5
 550 560 570 580 590 600
 ACGGAGCCGATTACCGACGCCGACCGACACCGAGGCCGCATTCCGCCGTAGGCGACAAAG
 isGlyAlaAspTyrArgArgProThrAspThrGluAlaAlaPheAlaValGlyAspLysV
 10
 610 620 630 640 650 660
 TCATCGTGCGGTCCGACGCCCTCACCGAACACCCACACCCGCCGCCGGATACGTCCGCG
 alIleValArgSerAspAlaSerProAsnThrHisThrArgArgAlaGlyTyrValArgG
 15
 670 680 690 700 710 720
 GTCGTGTCCGCGAAGTCGTGGCGACCCACGGCGCGTATGTCTTCCGGACACCAACGCAC
 lyArgValGlyGluValValAlaThrHisGlyAlaTyrValPheProAspThrAsnAlaL
 20
 730 740 750 760 770 780
 TCGGCGCCGCGGAAAGCCCCGAACACCTGTACACCGTGCGGTTCTCGGCGACCGAGTTGT
 euGlyAlaGlyGluSerProGluHisLeuTyrThrValArgPheSerAlaThrGluLeuT
 25
 790 800 810 820 830 840
 GGGGTGAACCTGCCGCCCGGAACGTCTCAATCACATCGACGTGTTCCGAACCGTATCTGC
 rpGlyGluProAlaAlaProAsnValValAsnHisIleAspValPheGluProTyrLeuL
 30
 850 860 870 880 890 900
 TACCGGCCTGACCAGGTCATCCGGTCCACCCAGCGAGACGTCCCTTCACCACAGACAGAA
 euProAla
 35
 910 920 930 940 950 960
 ACGAGCCACCCCGATGACCGCCCAATCCCGTCCAGGGCACGTTGCCACGATCGAACG
 MetThrAlaHisAsnProValGlnGlyThrLeuProArgSerAsnG
 Subunit α (L)
 40
 970 980 990 1000 1010 1020
 AGGAGATCGCCGCACGCGTGAAGGCCATGGAGGCCATCCTCGTCGACAAGGGCCTGATCT
 luGluIleAlaAlaArgValLysAlaMetGluAlaIleLeuValAspLysGlyLeuIleS
 45
 1030 1040 1050 1060 1070 1080
 CCACCGACGCCATCGACCACATGTCCTCGGTCTACGAGAACGAGGTCCGTCTCAACTCG
 erThrAspAlaIleAspHisMetSerSerValTyrGluAsnGluValGlyProGlnLeuG
 50
 1090 1100 1110 1120 1130 1140
 GCGCCAAGATCGTCGCCCGCGCCTGGGTTCGATCCCGAGTTCAAGCAGCGCCTGCTCACCG
 lyAlaLysIleValAlaArgAlaTrpValAspProGluPheLysGlnArgLeuLeuThrA
 55
 1150 1160 1170 1180 1190 1200
 ACGCCACCAGCGCCTGCCGTGAAATGGGCGTCCGGCGGCATGCAGGGCGAAGAAATGGTCC
 spAlaThrSerAlaCysArgGluMetGlyValGlyGlyMetGlnGlyGluGluMetValV
 60
 1210 1220 1230 1240 1250 1260
 TGCTGGAAAACACCGGCACGGTCCACAACATGGTTCGTATGTACCTTGTGCTCGTGCTATC
 alLeuGluAsnThrGlyThrValHisAsnMetValValCysThrLeuCysSerCysTyrP
 65
 1270 1280 1290 1300 1310 1320
 CGTGGCCGTTCTCGGCCTGCCACCCAACCTGGTACAAGTACCCCGCCTACCGCGCCCGCG
 roTrpProValLeuGlyLeuProProAsnTrpTyrLysTyrProAlaTyrArgAlaArgA
 70
 1330 1340 1350 1360 1370 1380
 CTGTCCGCGACCCCGAGGTGTGCTGGCCGAATTCGGATATACCCCGACCCGTGACGTCC
 laValArgAspProArgGlyValLeuAlaGluPheGlyTyrThrProAspProAspValG

1390 1400 1410 1420 1430 1440
 AGATCCGGATATGGGACTCGAGTGCCGAACCTTCGCTACTGGGTCCTGCCGCAACGCCACG
 5 luIleArgIleTrpAspSerSerAlaGluLeuArgTyrTrpValLeuProGlnArgProA
 1450 1460 1470 1480 1490 1500
 CCGGCACCGAGAACTTCACCGAAGAACAACCTCGCCGACCTCGTCACCCGCGACTCGCTCA
 10 laGlyThrGluAsnPheThrGluGluGlnLeuAlaAspLeuValThrArgAspSerLeuI
 1510 1520 1530 1540 1550 1560
 TCGGCGTATCCGTCCCCACCCACCCAGCAAGGCCTGACATGCCCCGACTCAACGAACAA
 15 leGlyValSerValProThrThrProSerLysAla
 1570 1580 1590 1600 1610 1620
 CCCCACCCGGGTCTCGAAGCCAACCTCGGCGACCTGGTACAGAATCTGCCGTTCAACGAA
 1630 1640 1650 1660 1670 1680
 CGAATCCCCCGCCGCTCCGGCGAGGTCGCCTTCGATCAGGCCTGGGAGATCCGCGCCTTC
 1690 1700 1710 1720 1730
 AGCATTGCCACCGCATTGCATGGCCAGGGCCGATTCTGAATGGGACGAATTC

Claims

1. A DNA^(H) fragment encoding a polypeptide having nitrile hydratase activity, said polypeptide comprising the $\alpha^{(H)}$ -subunit as defined in the Sequence Listing by SEQ ID: No. 1 and the $\beta^{(H)}$ -subunit as defined in the Sequence Listing by SEQ ID: No. 2.
2. A DNA^(L) fragment encoding a polypeptide having nitrile hydratase activity, said polypeptide comprising the $\alpha^{(L)}$ -subunit as defined in the Sequence Listing by SEQ ID: No. 3 and the $\beta^{(L)}$ -subunit as defined in the Sequence Listing by SEQ ID: No. 4.
3. The DNA^(H) fragment of claim 1 which contains the nucleotide sequences of the $\alpha^{(H)}$ -subunit as defined in the Sequence Listing by SEQ ID: No. 5 and the $\beta^{(H)}$ -subunit as defined in the Sequence Listing by SEQ ID: No. 6.
4. The DNA^(L) fragment of claim 2 which contains the nucleotide sequences of the $\alpha^{(L)}$ -subunit as defined in the Sequence Listing by SEQ ID: No. 7 and the $\beta^{(L)}$ -subunit as defined in the Sequence Listing by SEQ ID: No. 8.
5. A recombinant DNA comprising a DNA^(H) or DNA^(L) of any one of claims 1-4 in a vector.
6. A transformant transformed with the recombinant DNA of claim 5.
7. A method of producing nitrile hydratase which comprises culturing the transformant as claimed in claim 6 and recovering nitrile hydratase from the culture.
8. A method of producing amides which comprises hydrating nitriles using nitrile hydratase obtained from the culture of the transformant of claim 6.
9. A method of producing amides which comprises culturing the transformant as claimed in claim 6, and hydrating nitriles to amides using the resultant culture, isolated bacterial cells, treated matter thereof, or a fixed material thereof.

FIG. 1

